

Abstract Submitted to the  
International Conference on Strongly Correlated Electron Systems  
University of Michigan, Ann Arbor  
August 6-10, 2001

## **Calculation of exchange integrals in transition metals oxides using the LDA+ $U$ approach**

Alexander Yaresko<sup>1</sup>, Roland Hayn<sup>2</sup>, Alexander Perlov<sup>3</sup>, Helge Rosner<sup>4</sup>, Peter Thalmeier<sup>1</sup>

<sup>1</sup> *Max-Planck Institute for the Chemical Physics of Solids, D-01187 Dresden, Germany*

<sup>2</sup> *Institute for Solid State and Materials Research, D-01171 Dresden, Germany*

<sup>3</sup> *Max-Planck Institute for the Physics of Complex Systems, D-01187 Dresden, Germany*

<sup>4</sup> *Department of Physics, University of California, 95616 Davis, California, USA*

Theoretical determination of effective exchange coupling constants between the magnetic ions in transition metal oxides is a general and long standing problem. We propose a solution to the problem based on the mapping of the energy of noncollinear spin-spiral states calculated with the LDA+ $U$  method to the mean-field solutions of the effective spin Hamiltonian. The approach is applied to  $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ , which can be described with good accuracy by a 2D Heisenberg model with only nearest-neighbor exchange, and to  $\text{Ba}_2\text{Cu}_3\text{O}_4\text{Cl}_2$ , which has two coupled anti-ferromagnetic spin systems. The same method is also used to determine the exchange constants between  $\text{V}^{4+}$  spins in the charge ordered low temperature phase of  $\text{NaV}_2\text{O}_5$  and their dependence on the assumed kind of the charge ordering. The sensitivity of the calculated exchange constants to the strength of the on-site Coulomb repulsion  $U$  is analyzed.